

A note on univoque self-Sturmian numbers

J.-P. Allouche*
 CNRS, LRI
 Bâtiment 490
 Université Paris-Sud
 F-91405 Orsay Cedex
 France
 allouche@lri.fr

Abstract

We compare two sets of (infinite) binary sequences whose suffixes satisfy extremal conditions: one occurs when studying iterations of a unimodal continuous map from the unit interval into itself, but it also characterizes univoque real numbers; the other is an equivalent definition of characteristic Sturmian sequences. As a corollary to our study we obtain that a real number β in $(1, 2)$ is univoque and self-Sturmian if and only if the β -expansion of 1 is of the form $1v$, where v is a characteristic Sturmian sequence beginning itself in 1.

Keywords: Sturmian sequences, univoque numbers, self-Sturmian numbers.

MSC: 11A63, 68R15.

1 Introduction

The kneading sequences of a unimodal continuous map f from $[0, 1]$ into itself, with $f(1) = 0$ and $\sup f = 1$ are classically studied by first looking at the combinatorial properties of the kneading sequence of 1. Cosnard proved that, using a simple bijection on binary sequences (namely mapping the sequence $(x_n)_{n \geq 0}$ to $(y_n)_{n \geq 0}$, where $y_n := \sum_{0 \leq j \leq n} x_j \bmod 2$), the set of kneading sequences of 1 for all maps f as above, maps to the set Γ defined by

$$\Gamma := \{u = (u_n)_{n \geq 0} \in \{0, 1\}^{\mathbb{N}}, \forall k \geq 0, \bar{u} \leq S^k u \leq u\}$$

where $\bar{u} = (\bar{u}_n)_{n \geq 0}$ is the sequence defined by $\bar{u}_n := 1 - u_n$, where S^k is the k th iterate of the shift (i.e., $S^k((u_n)_{n \geq 0}) := (u_{n+k})_{n \geq 0}$), and where $\leq$ is the lexicographical order on sequences induced by $0 < 1$. (See [10, 2]; actually the relevant set there is $\Gamma \setminus \{(10)^\infty\}$. See also [1] for a detailed combinatorial study of the set Γ .)

*supported by MENESR, ACI NIM 154 Numération.

A slight modification of the set Γ describes the expansions of 1 in bases β , where $\beta \in (1, 2)$ is a *univoque* number, i.e., a number such that 1 admits only one expansion in base β (see [11, Remark 1, page 379], see also [4] and the bibliography therein): the set of all the expansions of 1 in bases $\beta \in (1, 2)$, where β is univoque, is the set

$$\Gamma_1 := \{u = (u_n)_{n \geq 0} \in \{0, 1\}^{\mathbb{N}}, \forall k \geq 0, \bar{u} < S^k u < u\}.$$

Remark 1 Note that a binary sequence belongs to Γ_1 if and only if it belongs to Γ and is not purely periodic.

Other sequences can be defined by extremal properties of their suffixes: characteristic Sturmian sequences and Sturmian sequences. More precisely the following results can be found in several papers (see in particular [15, 16, 8, 14, 7]; see also the survey [6] and the discussion therein).

A binary sequence $u = (u_n)_{n \geq 0}$ is characteristic Sturmian if and only if it is not periodic and belongs to the set Ξ , where

$$\Xi := \{u = (u_n)_{n \geq 0} \in \{0, 1\}^{\mathbb{N}}, \forall k \geq 0, 0u \leq S^k u \leq 1u\}.$$

A binary sequence $u = (u_n)_{n \geq 0}$ is Sturmian if and only if it is not periodic and there exists a binary sequence $v = (v_n)_{n \geq 0}$, such that u belongs to Ξ_v , where

$$\Xi_v := \{u = (u_n)_{n \geq 0} \in \{0, 1\}^{\mathbb{N}}, \forall k \geq 0, 0v \leq S^k u \leq 1v\}.$$

The sequence v has the property that $1v = \sup_k S^k u$ and $0v = \inf_k S^k u$. This is the characteristic Sturmian sequence having the same slope as u .

Remark 2 The reader can find everything on Sturmian sequences in [13, Chapter 2]. A hint for the proof of the two assertions above is that a sequence is Sturmian if and only if it is not periodic and for any binary (finite) word w , the words $0w0$ and $1w1$ cannot be simultaneously factors of the sequence; furthermore a sequence u is characteristic Sturmian if and only if $0u$ and $1u$ are both Sturmian.

2 Comparing the sets Γ and Ξ

The analogy between the definitions of Γ and Ξ suggests the natural question whether any sequence can belong to their intersection. The disappointing answer is the following proposition.

Proposition 1 *A sequence $u \in \{0, 1\}^{\mathbb{N}}$ belongs to $\Gamma \cap \Xi$ if and only if it is equal to 1^∞ or there exists $j \geq 1$ such that $u = (1^j 0)^\infty$.*

Proof. If the sequence u belongs to Γ , we have in particular $u \geq \bar{u}$. Hence $u = 1w$ for some binary sequence w . If u is not equal to 1^∞ (which clearly belongs to $\Gamma \cap \Xi$), let us write $u = 1^j 0z$ for some integer $j \geq 1$ and some binary sequence z . Since u belongs to Γ we have $S^{j+1}u \leq u$, i.e., $z \leq u$. Now u belongs to Ξ , thus $S^j u \geq 0u$, i.e., $0z \geq 0u$, hence $z \geq u$. This gives $z = u$. Hence $u = (1^j 0)^\infty$, which in turn clearly belongs to $\Gamma \cap \Xi$. $\square$

The next question is whether a Sturmian sequence can belong to Γ . The answer is more interesting.

Proposition 2 *A (binary) Sturmian sequence u belongs to Γ if and only if there exists a characteristic Sturmian sequence v such that v begins in 1 and $u = 1v$.*

Proof.

Let us first suppose that the Sturmian sequence u belongs to Γ . As above, since u belongs to Γ , u begins in 1. Hence $u = 1w$ for some binary sequence w . The inequalities $S^k u \leq u$ for all $k \geq 0$ imply that $\sup_k S^k u = u$ (the inequality $\geq$ is trivial since $S^0 u = u$). This can be written $\sup_k S^k u = 1w$. On the other hand u is Sturmian, hence there exists a binary sequence v such that u belongs to Ξ_v . We also know that v is such that $1v = \sup_{k \geq 0} S^k u$. Hence $v = w$. Now $\inf_k S^k u = 0v = 0w$. But $S^k u \geq \bar{u}$ for all $k \geq 0$, since u belongs to Γ . Hence $0v = 0w \geq \bar{u} = 0\bar{w}$, thus $w \geq \bar{w}$, hence w begins in 1.

If, conversely, $u = 1v$ where v is a characteristic Sturmian sequence (which actually implies that u is Sturmian) beginning in 1, we first note that $0v \leq S^k v \leq 1v$ for all $k \geq 0$. Hence, immediately, $S^k u \leq 1v = u$ for all $k \geq 0$ (using that $S^{k+1}u = S^k v$ and that $S^0 u = u = 1v$). And also $S^k u \geq 0v \geq 0\bar{v} = \bar{u}$ (using furthermore that $v \geq \bar{v}$ since v begins in 1, and that $S^0 u = u \geq \bar{u}$ since u begins in 1). $\square$

Remark 3 We see in particular that a Sturmian sequence belonging to Γ must begin in 11. This is not surprising since the only sequence belonging to Γ that begins in 10 is $(10)^\infty$. This is a particular case of a lemma in [1]: *if a sequence t belonging to Γ begins with $m\bar{m}$, where m is a (finite) nonempty binary word, then $t = (m\bar{m})^\infty$.*

3 Univoque self-Sturmian numbers

Several papers were devoted to univoque numbers having an extra property. For example:

- the smallest univoque number in $(1, 2)$ is determined in [12]; it is related to the celebrated Thue-Morse sequence and was proven transcendental in [3];
- a detailed study of univoque Pisot numbers belonging to $(1, 2)$ can be found in [5].

Self-Sturmian numbers were introduced in [9]: these are the real numbers β such that the greedy β -expansion of 1 is a Sturmian sequence on some two-digit alphabet. It is tempting to ask which univoque numbers are self-Sturmian. We restrict the study to the numbers in $(1, 2)$ for simplicity.

Proposition 3 *The real self-Sturmian numbers in $(1, 2)$ that are univoque are exactly the real numbers β such that $1 = \sum_{n \geq 1} \frac{u_n}{\beta^n}$, where $u = (u_n)_{n \geq 0}$ is a binary sequence of the form $u = 1v$, with v a characteristic Sturmian sequence beginning in 1.*

Proof. This is a rephrasing of Proposition 2. $\square$

Remark 4 - The equality $1 = \sum_{n \geq 1} \frac{u_n}{\beta^n}$, where $u = (u_n)_{n \geq 0}$ is a binary sequence, uniquely determines the real number β in $(1, 2)$.

- Self-Sturmian numbers correspond to Sturmian sequences of the form $u = 1v$, where v is any characteristic Sturmian sequence (see [9, Remark p. 399]). All self-Sturmian numbers are transcendental [9].

4 Acknowledgments

The author wants to thank Amy Glen for her comments on a previous version of this note.

References

- [1] J.-P. Allouche, *Théorie des nombres et automates*, Thèse d'État, Université Bordeaux I, 1983.
- [2] J.-P. Allouche, M. Cosnard, Itérations de fonctions unimodales et suites engendrées par automates, *C. R. Acad. Sci. Paris, Série I* **296** (1983) 159–162.
- [3] J.-P. Allouche, M. Cosnard, The Komornik-Loreti constant is transcendental, *Amer. Math. Monthly* **107** (2000) 448–449.
- [4] J.-P. Allouche, M. Cosnard, Non-integer bases, iteration of continuous real maps, and an arithmetic self-similar set, *Acta Math. Hung.* **91** (2001) 325–332.
- [5] J.-P. Allouche, C. Frougny, K. G. Hare, On univoque Pisot numbers, *Math. Comp.*, to appear.
- [6] J.-P. Allouche, A. Glen, Extremal properties of (epi)sturmian sequences and distribution modulo 1, Preprint (2007).
- [7] Y. Bugeaud, A. Dubickas, Fractional parts of powers and Sturmian words, *C. R. Math. Acad. Sci. Paris* **341** (2005) 69–74.
- [8] S. Bullett, P. Sentenac, Ordered orbits of the shift, square roots, and the devil's staircase, *Math. Proc. Cambridge Philos. Soc.* **115** (1994) 451–481.
- [9] D. P. Chi, D. Kwon, Sturmian words, β -shifts, and transcendence, *Theoret. Comput. Sci.* **321** (2004) 395–404.

- [10] M. Cosnard, Étude de la classification topologique des fonctions unimodales, *Ann. Inst. Fourier* **35** (1985) 59–77.
- [11] P. Erdős, I. Joó, V. Komornik, Characterization of the unique expansions $1 = \sum q^{-n_i}$ and related problems, *Bull. Soc. Math. France* **118** (1990) 377–390.
- [12] V. Komornik, P. Loreti, Unique developments in non-integer bases, *Amer. Math. Monthly* **105** (1998) 636–639.
- [13] M. Lothaire, *Algebraic Combinatorics On Words, Encyclopedia of Mathematics and its Applications*, vol. 90, Cambridge University Press, 2002.
- [14] G. Pirillo, Inequalities characterizing standard Sturmian words, *Pure Math. Appl.* **14** (2003) 141–144.
- [15] P. Veerman, Symbolic dynamics and rotation numbers, *Physica A* **134** (1986) 543–576.
- [16] P. Veerman, Symbolic dynamics of order-preserving orbits, *Physica D* **29** (1987) 191–201.